

Utah State University

DigitalCommons@USU

Bu

Bee Lab

10-1-1901

Description of a New Genus of Bees from India

Peter Cameron

Follow this and additional works at: https://digitalcommons.usu.edu/bee_lab_bu



Part of the [Entomology Commons](#)

Recommended Citation

Cameron, Peter, "Description of a New Genus of Bees from India" (1901). *Bu*. Paper 179.
https://digitalcommons.usu.edu/bee_lab_bu/179

This Article is brought to you for free and open access by the Bee Lab at DigitalCommons@USU. It has been accepted for inclusion in Bu by an authorized administrator of DigitalCommons@USU. For more information, please contact digitalcommons@usu.edu.



DESCRIPTION OF A NEW GENUS OF BEES FROM INDIA.

By PETER CAMERON.

AGLAOPIS, gen. nov.

Wings short, not reaching to the apex of the third abdominal segment. There are two closed cubital cellules; the first is longer than the second above and below; the radial cellule has the apex rounded and not much narrowed; the first recurrent nervure is interstitial; the second is received not far from the apex of the cellule. Eyes large, slightly converging below, and reaching to the base of the mandibles; the ocelli are in a curve. Labrum visible, short, obliquely depressed. Mandibles bidentate; the apical tooth is much larger than the other. Scutellum bluntly projecting laterally at the apex; the middle slightly projects, and is keeled; the post-scutellum is keeled in the middle. Median segment short; the basal area large. Legs short and stout, pilose; the tibiae and tarsi not densely covered with long hair. The basal segment of the abdomen is transverse at the base, and is bordered at the base by a distinct keel; the apical segment is longer than the penultimate, and is distinctly incised at the apex; the sides of the incision are straight and oblique, and form two distinct lobes, which become gradually narrowed from the base to the apex; the lower edge is stoutly keeled. The antennae are short and stout; the third joint is not much narrowed; the head is well-developed behind the eyes, with the occiput transverse; the apex of the scutellum does not project over the post-scutellum, and only over the median segment at the sides; the claws and spurs are simple; the tegulae are large; the head and thorax are thickly covered with short white pubescence; the abdominal segments are narrowly banded at the apex with white pubescence. There is no ventral scopa. The front calcaria are normally curved; the claws simple.

Belongs, if anywhere, to the Stelididae, and comes nearest perhaps to *Parevaspis*, which may be known from it by the projecting apex of the scutellum, which is incised at the apex; by the second recurrent nervure being received beyond the second transverse cubital nervure, and not in the second cubital cellule; by the apex of the abdomen not being deeply incised in the middle in the female; and by the longer wings. The wings in the present genus are shorter than usual. It is easily known by the form of the scutellum, by the transversely keeled basal segment of the abdomen, and by the deep incision in the apical segment. The colour—black, with the basal two or three segments of the abdomen red—is peculiar, and does not occur with any other Indian species. The genus is doubtless, like *Stelis* and *Parevaspis*, a parasitic one.

AGLAOPIS BREVIPENNIS, sp. nov.

Black, thickly covered with white pubescence; the apices of the abdominal segments banded with white pubescence; the greater part

of the f
third, re
Long. 6

Hal

Ante

the scap
closely p
thickly
vertex m
black, r
white pu
tellum of
which is
scutellum
of the m
slope lat
thickly c
with whit
distinctly
stronger

BUTTER

THE

Austria a
from Jun
follows :-
26th to 2
Buda Pe
Orsowa,

At Sa
almost e
east of t

At Be
there is a
nately th
day—Jun

At M
the best
lucilla wa

At Bu
were mos
was mos
and-pini
covered
slopes.

NEW GENUS OF BEES FROM INDIA.

BY PETER CAMERON.

LALAPIS, gen. nov.

ing to the apex of the third abdominal seg-
ed cubital cellules; the first is longer than
y; the radial cellule has the apex rounded
the first recurrent nervure is interstitial; the
rom the apex of the cellule. Eyes large,
and reaching to the base of the mandibles;
Labrum visible, short, obliquely depressed.
apical tooth is much larger than the other.
g laterally at the apex; the middle slightly
the post-scutellum is keeled in the middle.
basal area large. Legs short and stout,
not densely covered with long hair. The
en is transverse at the base, and is bordered
el; the apical segment is longer than the
tly incised at the apex; the sides of the
lique, and form two distinct lobes, which
from the base to the apex; the lower edge
ennæ are short and stout; the third joint
e head is well-developed behind the eyes,
the apex of the scutellum does not pro-
and only over the median segment at the
s are simple; the tegulæ are large; the
covered with short white pubescence; the
arrowly banded at the apex with white
ventral scopa. The front calcaria are
simple.

to the Stelididæ, and comes nearest
ich may be known from it by the pro-
um, which is incised at the apex; by
ure being received beyond the second
, and not in the second cubital cellule;
men not being deeply incised in the
by the longer wings. The wings in
ter than usual. It is easily known by
m, by the transversely keeled basal
nd by the deep incision in the apical
ack, with the basal two or three seg-
—is peculiar, and does not occur with
The genus is doubtless, like *Stelis* and

BREVIPENNIS, sp. nov.

ith white pubescence; the apices of the
with white pubescence; the greater part

of the first and the whole of the second segment, and sometimes the
third, red; the wings hyaline, the stigma and nervures black. Female.
Long. 6-7 mm.

Hab. Bombay (G. A. J. Rothney).

Antennæ black, the flagellum with a faint brownish tint; bare,
the scape covered with white hair. Front and vertex strongly and
closely punctured; the front, the sides of the face, and of the clypeus
thickly covered with white pubescence; the central parts and the
vertex much more sparsely covered with similar pubescence. Mandibles
black, rufous near the apex; the base punctured, and covered with
white pubescence; the middle above hollowed. Mesonotum and scu-
tellum closely rugosely punctured, and covered with white pubescence,
which is thicker and more fulvous in tint round the eyes; the post-
scutellum is thickly covered with white longish pubescence. The base
of the median segment is stoutly longitudinally striated; its apical
slope laterally is covered thickly with white pubescence. Pleuræ
thickly covered with white pubescence. Legs black, thickly covered
with white pubescence; the calcaria testaceous. Abdomen closely and
distinctly punctured; the punctuation on the basal two segments is
stronger and more widely separated than it is on the others.

BUTTERFLY COLLECTING IN AUSTRO-HUNGARY IN 1900.

BY HENRY C. LANG, M.D., F.E.S.

THE following notes are a record of butterfly collecting in
Austria and Hungary during one month of the summer of 1900,
from June 21st to July 21st. The dates and localities were as
follows:—Salzburg, June 21st to 28th; Berchtesgaden, June
26th to 28th; Modling, near Vienna, June 29th to July 2nd;
Buda Pesth, July 3rd to 9th; Herculesbad, July 12th to 20th;
Orsowa, July 20th.

At Salzburg there were very few butterflies on the low ground;
almost everything was collected on the wooded hills a few miles
east of the town.

At Berchtesgaden, in Bavaria, but a few miles from Salzburg,
there is a fine opportunity for mountain collecting, but unfortu-
nately the weather was dull or rainy, with the exception of one
day—June 27th—when I collected in the woods above the village.

At Modling, near Vienna, the weather was fine, but some of
the best species found there were not yet on the wing (one *Neptis*
lucilla was taken).

At Buda Pesth we had very unfavourable weather; the days
were mostly showery, and at times windy and cold. Collecting
was mostly on the Schwabenberg, a mountain reached by rack-
and-pinion railway from the town. It is an interesting locality,
covered with woods of oak and beech, and with open grassy
slopes.